

SHANGHAI JIAO TONG UNIVERSITY
LECTURE 2
2017

Anthony J. Leggett

Department of Physics

University of Illinois at Urbana-Champaign, USA

and

Director, Center for Complex Physics

Shanghai Jiao Tong University



Reminder: the vector potential $\mathbf{A}(\mathbf{r}, t)$ in classical mechanics

The equation of motion of a charged particle in an electric field \mathbf{E} and magnetic field \mathbf{B} :

$$m \frac{d\mathbf{v}}{dt} = e(\mathbf{E} + \mathbf{v} \times \mathbf{B}) (+\mathbf{F}_{\text{non-em}})$$

How to find $H(\mathbf{r}, \mathbf{p})$ s.t.

$$\frac{d\mathbf{r}}{dt} = \frac{\partial H}{\partial \mathbf{p}}, \quad \frac{d\mathbf{p}}{dt} = -\frac{\partial H}{\partial \mathbf{r}} \quad ?$$

Solution: define $A(\mathbf{r}, t)$ s.t.

$$\mathbf{E}(\mathbf{r}, t) = -\frac{\partial \mathbf{A}(\mathbf{r}, t)}{\partial t}, \quad \mathbf{B}(\mathbf{r}, t) = \nabla \times \mathbf{A}(\mathbf{r}, t)$$

(hence $\nabla \times \mathbf{E} = \partial \mathbf{B} / \partial t$)

and put

↑
Faraday

$$H(\mathbf{r}, \mathbf{p}) = (\mathbf{p} - e\mathbf{A}(\mathbf{r}, t))^2 / 2m \quad (+V_{\text{non-em}})$$

This works! [Recall:

$$\frac{d\mathbf{A}(\mathbf{r}, t)}{dt} = \frac{\partial \mathbf{A}(\mathbf{r}, t)}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{A}, \quad (\mathbf{v} \times (\nabla \times \mathbf{A}))_i = v_j \partial_i A_j - (\mathbf{v} \cdot \nabla) A_i]$$

$$\text{note: } \mathbf{v} \equiv \frac{d\mathbf{r}}{dt} = \frac{\partial H}{\partial \mathbf{p}} = \frac{1}{m} (\mathbf{p} - e\mathbf{A}(\mathbf{r}, t)) \quad (\neq \mathbf{p}/m)$$

so $H(\mathbf{r}, \mathbf{p}) = \frac{1}{2} m \mathbf{v}^2 =$ kinetic energy (only)! (but expressed in terms of \mathbf{p} and \mathbf{A}).

Quantum mechanics:

$\mathbf{p} \rightarrow -i\hbar\nabla$ so KE is

$$\hat{H}_K = (-i\hbar\nabla - e\mathbf{A})^2/2m$$

and so, including possible $V_{non-em} \equiv V(\mathbf{r})$,

$$\hat{H} = \frac{1}{2m} (-i\hbar\nabla - e\mathbf{A}(rt))^2 + V(\mathbf{r})$$

In CM (classical mechanics), all effects obtainable from $\mathbf{A}(\mathbf{r}t)$ are equally derivable only from $\mathbf{E}(\mathbf{r}t)$ and $\mathbf{B}(\mathbf{r}t) \Rightarrow$ vector potential redundant. In QM (quantum mechanics) this is not true: $\mathbf{A}(\mathbf{r}t)$ has a “life of its own”!


Single charged particle on thin ring

If field \mathbf{B} through ring is uniform, can take

$$\mathbf{A} \equiv A_\theta \hat{\boldsymbol{\theta}}, A_\theta \equiv \frac{1}{2}BR$$

then flux Φ through ring is

$$\Phi \equiv \pi R^2 B \Rightarrow A_\theta = \Phi/2\pi R$$



TISE for $\psi \equiv \psi(\theta)$ is

$$\hat{H}\psi(\theta) \equiv \frac{1}{2m} \left(-i\hbar \nabla - e\mathbf{A}(\mathbf{r}) \right)^2 \psi(\theta) = E\psi(\theta)$$

only nonzero component of ∇ is $\nabla_\theta = \frac{1}{R} \frac{\partial}{\partial \theta}$

and only component of \mathbf{A} is A_θ , so

$$\frac{\hbar^2}{2mR^2} \left(-i \frac{\partial}{\partial \theta} - \frac{e}{\hbar} A_\theta R \right)^2 \psi(\theta) = E\psi(\theta)$$

or putting $A_\theta = \Phi/2\pi R$ and defining $\Phi_o^{sp} \equiv h/e$

(single-particle)
flux quantum

$$\frac{\hbar^2}{2mR^2} \left(-i \frac{\partial}{\partial \theta} - \Phi/\Phi_o^{sp} \right)^2 \psi(\theta) = E\psi(\theta)$$


$$= \hat{L}_z \text{ (angular momentum in units of } \hbar)$$

Formal solution is

$$\psi(\theta) = \exp ik\theta, \quad (k \text{ arbitrary}), \quad E = \frac{\hbar^2}{2mR^2} \left(k - \Phi/\Phi_o^{sp} \right)^2$$

However, crucial point:

$\psi(\theta)$ must be single-valued, i.e. $\psi(\theta + 2n\pi) = \psi(\theta)$ (SVBC)

single-valuedness
boundary condition 

Hence, only allowed values of k are integers $\ell = 0, \pm 1, \pm 2 \dots$

(i.e. angular momentum \hat{L}_z is quantized in units of \hbar)

Thus,

$$\psi_\ell(\theta) = \exp i\ell\theta, \quad \ell = 0, \pm 1, \pm 2,$$

$$E_\ell = \frac{\hbar^2}{2mR^2} (\ell - \Phi/\Phi_o^{sp})^2$$

$$j_\ell = \frac{e\hbar}{m/R} (\ell - \Phi/\Phi_o^{sp})$$

For $\Phi < \Phi_o^{sp}/2$,

$$\text{GS has } \ell = 0 \Rightarrow p_\theta \equiv L_z/R = \ell\hbar/R = 0$$

However, recall:

$$\mathbf{v} = (\mathbf{p} - e\mathbf{A})/m \Rightarrow \mathbf{v}_\theta = -eA_\theta/m$$

$\Rightarrow \mathbf{j}_\theta \equiv e\mathbf{v}_\theta = -(e^2/m)A_\theta \neq 0$ in general, in
sense to produce magnetic field opposite to
 $\mathbf{B} \Rightarrow$ GS is diamagnetic,

$$\mathbf{j}_\theta = -(e^2/m)A_\theta \neq 0$$




Single charged particle on ring: two notes

1. What is the situation in classical mechanics?

We can still formally introduce $\ell \equiv L_z/\hbar$ and write

$$E(\ell) = \frac{\hbar^2}{2mR^2} \left(\ell - \Phi/\Phi_0^{sp} \right)^2$$

but now there is no restriction on ℓ (SVBC is meaningless since no wave function!) so now GS always corresponds to $\ell = \Phi/\Phi_0^{sp}$, equivalent to $\mathbf{j}_\theta = 0$ (no diamagnetism).



However, consider **time-dependent** problem: since motion is restricted to ring, Lorenz force $\mathbf{v} \times \mathbf{B}$ is irrelevant and we have by Newton II


$$m \frac{d\mathbf{v}}{dt} = \mathbf{E}(t) = -\frac{\partial \mathbf{A}}{\partial t}$$

or

$$m \frac{dv_\theta}{dt} = -\frac{dA_\theta}{dt}$$

If at $t = 0$ $v_\theta = 0$,
solution is simply

$$v_\theta(t) = -(e/m)A_\theta(t)$$




and in particular for $t = t_0$

$$v_\theta = -(e/m)A_f \Rightarrow j_\theta = -(e^2/m)A_f$$

as in Quantum Mechanics case. However, this is not the lowest-energy state, so scattering by walls, etc. will reduce j_θ to zero.

2. Aharonov-Bohm effect

note induced diamagnetic current depends only on **total trapped flux Φ** , not on details of how it is produced. Hence in particular can get nonzero effect even when **$B = 0$ everywhere on ring!** (e.g. B produced by “Helmholtz coil”)



Atomic diamagnetism

To the extent that argument applies, velocity of electrons at radius r given by

$$\mathbf{v}(r) = -e\mathbf{A}(r)/m$$

but electric current density $\mathbf{j}(r) = n(r)e\mathbf{v}(r)$, hence

$$\mathbf{j}(r) = \frac{-n(r)e^2}{m}\mathbf{A}(r)$$

Circulating current produces magnetic field ΔB **opposite** to original one.

Estimate order of magnitude: take

$$A \sim BR_{at}, J \sim R_{at}^2 j$$

$$\sim R_{at}^2 n e^2 A / m \sim (N e^2 / m R_{at}) A \sim (N e^2 / m) B,$$

$$\Delta B \sim \frac{\mu_0 J}{R_{at}} \sim \left(\frac{N e^2 \mu_0}{m R_{at}} \right) B:$$

$(N e^2 \mu_0 / m R_{at}) \sim 10^{-6}$, so effect very small.

Superconductors: London phenomenology

Basic postulate: as in atomic diamagnetism,

$$\mathbf{j}(\mathbf{r}) = \frac{-ne^2}{m} \mathbf{A}(\mathbf{r})$$

Combine with Maxwell's equation

$$\mathbf{j} = \nabla \times \mathbf{H} \equiv \frac{1}{\mu_0} \nabla \times \mathbf{B} = \frac{1}{\mu_0} \nabla \times (\nabla \times \mathbf{A}) = -\frac{1}{\mu_0} \nabla^2 \mathbf{A} \quad (\text{div } \mathbf{A} = 0)$$

gives

$$\nabla^2 \mathbf{A} = +\frac{ne^2}{m} \mu_0 \mathbf{A}$$

or taking curl.

$$\nabla^2 \mathbf{B} = \frac{ne^2 \mu_0}{m} \mathbf{B} \equiv \lambda_L^{-2} \mathbf{B}$$

London penetration depth

Hence, both in atomic diamagnets and in superconductors,

$$B \sim B_0 \exp -z/\lambda_L \quad (n(r), \text{ hence } \lambda_L, \text{ comparable in two cases})$$

Qualitative difference: in both cases $\lambda_L \sim 10^{-5}$ cm, but:

in atomic diamagnets, $\lambda_L \gg$ atomic size \Rightarrow effect very small

in superconductors, $\lambda_L \ll$ size of sample \Rightarrow effect spectacular: magnetic field totally excluded from bulk (Meissner effect)

Problems with London phenomenology

- A. Meissner effect is **thermodynamically stable** phenomenon, circulating supercurrents on **metastable**. Hence London argument does **not** explain stability of supercurrents! (↑: beware misleading statements in literature)
- B. No explanation of vanishing Peltier coefficient.
- C. Why do not **all** metals show Meissner effect?

Let's turn question C around: when does Meissner effect **not** occur?

1. Classical systems:

no restriction on $v_\theta \equiv (p_\theta - eA_\theta)/m$, and by Maxwellian statistical mechanics $P(v_\theta) \propto \exp\frac{1}{2}mv_\theta^2/kT$, hence from symmetry $\bar{v} = 0 \Rightarrow$ no circulating current \Rightarrow no diamagnetism (**Bohr-van Leeuwen theorem**)

2. Quantum Mechanics, but noninteracting particles obeying classical statistics: now p (or angular momentum L) is quantized

$$L = \ell \hbar, \quad \ell = \dots - 2, -1, 0, 1, 2 \dots$$

$$\text{and energy} \propto \left(\ell - \Phi / \Phi_o^{sp} \right)^2 \hbar^2 / 2mR^2 \quad \Phi_o^{sp} \equiv h/e$$

so

$$P(\ell) \propto \exp - \left\{ \left(\ell - \Phi / \Phi_o^{sp} \right)^2 \cdot \hbar^2 / 2mR^2 k_B T \right\} \quad (\Phi \lesssim \Phi_o^{sp})$$

Crucial point: under normal circumstances $k_B T \gg \hbar^2 / 2mR^2$, so can effectively replace discrete values of ℓ by continuum \Rightarrow back to classical mechanics.*

3. So, will only get Meissner effect if for some reason **all or most particles forced to be in same state**. Then the probability of angular momentum ℓ for this state is

$$P(\ell) \propto \exp - N_o \left(\ell - \Phi / \Phi_o \right)^2 \hbar^2 / 2mR^2 k_B T$$



Number of particles in same state

and provided $k_B T$, though $\gg \hbar^2 / 2mR^2$, is $\ll N \hbar^2 / 2mR^2$, can get results similar to atomic diamagnetism.

Does this ever happen? Yes, e.g. for **noninteracting gas of bosons!**

*Doesn't work for atomic diamagnetism because $\hbar^2 / 2mR^2$ is $\sim eV$,



hence $\gg k_B T$.

Electron volts

Summary of lecture 2:

- (1) In presence of electromagnetic vector potential $\mathbf{A}(\mathbf{r})$, Hamiltonian for single particle of charge e is

$$\hat{H} = p \left(\frac{-i\hbar\nabla - e\mathbf{A}(\mathbf{r})}{2m} \right)^2 + V(\mathbf{r})$$

- (2) For single particle on ring, in flux $\Phi < \frac{1}{2}h/e$, this leads in ground state to $j_\theta = -(e^2/m)A_\theta$

- (3) For a closed-shell atom, similar argument leads to

$$\mathbf{j}(\mathbf{r}) = -n \frac{(\mathbf{r})}{m} e^2 \mathbf{A}(\mathbf{r}) \quad (\text{diamagnetism}) \quad (*)$$

- (4) London phenomenology: assume (*) also describes superconductor

\Rightarrow Meissner effect

- (5) Difficulty: doesn't work for classical systems (Bohr – van Leeuwen theorem) nor (for $kT \gtrsim \hbar^2/mR^2$) for quantum systems obeying Maxwell-Boltzmann statistics

- (6) Difficulty can be overcome if for some reason

all particles forced to behave in same way.