

SHANGHAI JIAO TONG UNIVERSITY

LECTURE 4

2017

Anthony J. Leggett

Department of Physics

University of Illinois at Urbana-Champaign, USA

and

Director, Center for Complex Physics

Shanghai Jiao Tong University



Ginzburg – Landau (GL) theory of superconductivity

Formulated 1950 (pre – BCS): even in 2017 mostly adequate for “engineering” applications of superconductivity. (SC)

historically, takes its origins in general Landau-Lifshitz theory (1936) of 2nd order phase transitions, hence quantitative validity confined to T close to T_c . Here, I will start by arguing “with hindsight” for its qualitative validity at $T = 0$, and only later generalize to $T \neq 0$ and in particular $T \rightarrow T_c$.

Recall: could explain 3 major characteristics of SC state (persistent currents, Meissner effect, vanishing Peltier coefficient) by scenario in which fermionic pairs form effective bosons, and these undergo BEC. Suppose that N fermions form $N/2$ bosons, which are then condensed into **the same** 2 – particle state, Neglect for now relative wave function and df.

COM wave function of condensed pairs $\equiv \varphi(\mathbf{r})$



COM coordinate

What is energy E_o (at $T = 0$) of pairs as function(al) of $\varphi(\mathbf{r})$?

At first sight, for a single pair,

$$E_o\{\varphi(\mathbf{r})\} = \underbrace{-E_b}_{\text{binding energy of pair}} \int |\varphi(\mathbf{r})|^2 d\mathbf{r} + \frac{\hbar^2}{2m} \int \left| \left(\nabla - \frac{2ie}{\hbar} \mathbf{A}(\mathbf{r}) \right) \varphi(\mathbf{r}) \right|^2 d\mathbf{r}$$

2 electrons involved!

so, convenient to define

“order parameter” (OP)

$$\Psi(\mathbf{r}) \equiv \sqrt{\frac{N}{2}} \varphi(\mathbf{r})$$

$$\alpha_{T=0} \equiv E_b$$

then energy of $N/2$ pairs is

$$E_o\{\Psi(\mathbf{r})\} = \text{const.} + \int d\mathbf{r} \left\{ -\alpha_{T=0} |\Psi(\mathbf{r})|^2 + \frac{\hbar^2}{2m} \left| \left(\nabla - \frac{2ie}{\hbar} \mathbf{A}(\mathbf{r}) \right) \Psi(\mathbf{r}) \right|^2 \right\}$$

However, by itself this will not generate stability of supercurrents (lecture 3). To do so, must add (e.g.) term in $|\Psi|^4$... Adding also EM field term:

$$E_o\{\Psi(\mathbf{r})\} =$$

$$\int d\mathbf{r} \left\{ -\alpha_{T=0} |\Psi(\mathbf{r})|^2 + \frac{1}{2} \beta_{T=0} |\Psi(\mathbf{r})|^4 + \frac{\hbar^2}{2m} \left| \left(\nabla - \frac{2ie}{\hbar} \mathbf{A}(\mathbf{r}) \right) \Psi(\mathbf{r}) \right|^2 + \frac{1}{2} \mu_0^{-1} (\nabla \times \mathbf{A}(\mathbf{r}))^2 \right\}$$

“Standard” form of GL (free)
energy (at $T=0$)



Notes:

1. Normalization of $\Psi(r)$ is conventional and arbitrary. Rescaling

$$\Psi^1(r) \equiv q\Psi(r), \alpha'_{T=0} \equiv q^{-2}\alpha_{T=0}, \beta'_{T=0} \equiv q^{-4}\beta_{T=0}, \hbar^2/2m \rightarrow \hbar^2/2mq^2$$

leaves E_0 unchanged.

2. If $A(r) = 0$ and $\Psi(r) = \text{constant} \equiv \Psi_{T=0}^{eq}$, then value is

$$\Psi_{T=0}^{eq} = (\alpha_{T=0}/\beta_{T=0})^{1/2}$$

and energy is $E_{T=0}^{eq} = -\alpha_{T=0}^2/2\beta_{T=0}$

3. Minimization with respect to $A(r)$ ($\delta E_0/\delta A(r) = 0$) yields Maxwell's equation

$$\nabla \times \mathbf{H} = \mathbf{j}(r)$$

provided that we identify

$$\mathbf{j}(r) = \frac{e}{m} (\Psi^*(r)(-i\hbar\nabla - 2e\mathbf{A})\Psi(r) + c.c.)$$

4. Minimization with respect to $\Psi(r)$ ($\delta E_0/\delta\Psi(r) = 0$) yields

$$-\alpha_{T=0}\Psi(r) + \beta_{T=0}|\Psi(r)|^2\Psi(r) - \frac{\hbar^2}{2m} \left(\nabla - 2\frac{ie}{\hbar}A(r) \right)^2 \Psi(r) = 0$$

Note that for $A(r) = 0$ this defines a characteristic length

$$\xi_{T=0} \equiv \left(\frac{\hbar^2}{2m\alpha_{T=0}} \right)^{1/2}$$

A second characteristic length follows if we put $\Psi(r) = \text{constant} \equiv \Psi_0$ and compare terms in A^2 and $(\nabla \times A)^2$.

$$\lambda_{T=0} \equiv (2m/e^2\mu_0|\Psi_0|^2)^{-1/2}$$

or since with our normalization $\Psi_0 = \sqrt{N/2V}$,

$$\lambda_{T=0} \equiv (m/\mu_0 e^2 n)^{-1/2} \quad (\equiv \text{London penetration depth at } T = 0)$$

 electron density

Generalization to $T \neq 0$:

Free energy

All we need do is to let $E_0 \rightarrow F(T)$ and let the coefficients α_0, β_0 be T – dependent:

$$F(\Psi(r); T) = F_0(T) + \int dr \mathcal{F}(r, T)$$

$\mathcal{F}(r, T) \equiv$

$$-\alpha(T)|\Psi(r)|^2 + \frac{1}{2}\beta(T)|\Psi(r)|^4 + \frac{\hbar^2}{2m} \left| \left(\nabla - \frac{2ie}{\hbar} A(r) \right) \Psi(r) \right|^2 + \mu_0^{-1} (\nabla \times A(r))^2$$

standard form of GL free energy density

Note that there is now a contribution to $\alpha(T)$ (and possibly also $\beta(T)$) from the entropy term $-TS\{\Psi(r)\}$. (condensate itself carries no entropy, but electrons “liberated” from it do!)

Since $\Psi(\mathbf{r})$ describes S state, should $\rightarrow 0$ at some T_c , then $\alpha(T)$ should change sign there. Most natural choice (corresponding to $E_b \sim \text{constant}$, $S\{\Psi(r)\} \sim \text{constant} - \text{constant} |\Psi|^2$):

$$\alpha(T) = \alpha_0(T_c - T)$$

$$\beta(T) = \beta_0$$

For $T \neq 0$, all the above results go through with $\alpha_{T=0} \rightarrow \alpha(T)$, $\beta_{T=0} \rightarrow \beta(T)$. In particular in limit $T \rightarrow T_c$ from below:

$$\Psi_{eq}(T) = (\alpha(T)/\beta(T))^{1/2} = (\alpha_0/\beta_0)^{1/2}(T_c - T)^{1/2}$$

$$F_{eq}(T) - F_0(T) = -\alpha^2(T)/2\beta(T) = -(\alpha_0^2/2\beta_0)(T_c - T)^2$$



$$\xi(T) \propto [\alpha(T)]^{-1/2} \propto (T_c - T)^{-1/2}$$

$$\lambda(T) \propto [\Psi_{eq} \cdot (T)]^{-1} \propto (T_c - T)^{-1/2}$$

so ratio $\kappa \equiv \lambda/\xi$ is independent of T in limit $T \rightarrow T_c$. From $F_{eq}(T) - F_0(T) \propto (T_c - T)^2$, entropy S has no discontinuity at T_c ,

but sp. ht. $C_v \equiv TdS/dT$ has discontinuity

$T_c \alpha_0^2/\beta_0 \Rightarrow$ second order phase transition at T_c .




Some simple applications of the GL theory

A. Zero magnetic field ($\mathbf{A} = 0$)

1. Recovery of OP near wall (etc.)

Suppose boundary condition at $z = 0$ is that $\Psi \rightarrow 0$ (e.g. wall ferromagnetic).

Then must solve GL eq



$$\alpha(T)\Psi(z) + \beta_0|\Psi(z)|^2\Psi(z) - \frac{\hbar^2}{2m} \frac{\partial^2\Psi(z)}{\partial z^2} = 0$$

subject to boundary conditions

$$\Psi(z = 0) = 0$$

$$\Psi(z \rightarrow \infty) = \Psi_\infty \equiv (\alpha(T)/\beta_0)^{1/2}$$

Solution:


$$\Psi(z) = \Psi_\infty \tanh(z/2\xi(T))$$

$$\xi(T) \equiv (\hbar^2/2m\alpha(T))^{-1/2} \propto (T_c - T)^{-1/2}$$

Note: Ψ does **not** have to vanish at boundary with vacuum or insulator! [except on scale $\sim k_F^{-1}$ where N state wave functions do too] Thus, no objection to SC occurring in grains of dimension $\ll \xi(T)$. However, contact with N metal tends to suppress SC.

2. Current – carrying state in thin wire

If $d \ll \lambda(T)$, can neglect \mathbf{A} to first approximation



By symmetry,

$$\Psi = |\Psi| \exp i\varphi, |\Psi| = \text{constant}, \mathbf{j} = \frac{2e\hbar}{m} |\Psi|^2 \nabla\varphi$$

$$F = -\alpha(T)|\Psi|^2 + \frac{\beta_0}{2}|\Psi|^4 + \frac{\hbar^2}{2m}|\Psi|^2(\nabla\varphi)^2$$

“Superfluid velocity”
 $\left(\mathbf{v}_s = \frac{\hbar}{2m} \nabla\varphi \right)$

and we need to minimize this with respect to $|\Psi|$ for fixed $\nabla\varphi$.

Result:

$$|\Psi| = \left(\frac{\alpha(T) - \frac{\hbar^2}{2m} |\nabla\varphi|^2}{\beta_0} \right)^{1/2} \equiv \Psi_\infty (1 - \xi^2(T)) (\nabla\varphi)^2)^{1/2}$$

$\Rightarrow |\Psi| \rightarrow 0$ for $\nabla\varphi = \xi^{-1}(T)$ (condensation energy changes sign).

However, j is **nonmonotonic** function of $\nabla\varphi$, with maximum at point when $\nabla\varphi = \frac{1}{\sqrt{3}}\xi^{-1}(T)$. (at which point $|\Psi| = \sqrt{2/3}\Psi_\infty$).

Thus, critical current j_c given by


$$j_c = \frac{2e\hbar}{m} \cdot \frac{2}{3} \frac{\alpha(T)}{\beta_0} \frac{1}{\sqrt{3}} \xi^{-1}(T) \propto \alpha^{-1}(T) \xi^{-1}(T) \propto (1 - T/T_c)^{3/2}$$

B. Behavior in magnetic field

Because of the Meissner effect, any superconductor will completely expel a **weak** magnetic field. If we consider just the competition between the resulting state and the N state, we have (per unit volume)

$$\Delta E_{magn} = +\frac{1}{2}\mu_0 H^2$$

$$\Delta E_{cond} = -\alpha^2(T)/2\beta_0$$



and so
 (“thermodynamic”) critical field H_c is given by

$$H_c(T) = (\alpha^2(T)/\mu_0\beta_0)^{1/2} \propto (1 - T/T_c)$$


However, in general this works only for “type-I” superconductors in form of long cylinder parallel to field.

More generally:

- (a) in type-I superconductors, “intermediate” state forms with interleaved macroscopic regions of N and S .
- (b) in type-II superconductors, magnetic field “punches through” sample in the form of **vortex lines**.

Isolated vortex line

Consider $\lambda(T) \gg \xi(T)$
 (“extreme type-II”)



Magnetic field turns region
 $\sim \xi^2(T)$ normal, and “punches
 through” there. Field is screened
 out of bulk on scale $\lambda(T) \gg \xi(T)$

At distances $\gg \lambda(T)$, no current flows: however since

$$\mathbf{j} \propto \nabla\varphi - \frac{2e}{\hbar} \mathbf{A}$$

$\nabla\varphi$ and \mathbf{A} may be individually non zero, with $\nabla\varphi = \frac{2e}{\hbar} \mathbf{A}$.

But φ must be single-valued mod. 2π , hence

$$\oint \nabla\varphi \cdot d\ell = 2n\pi \Rightarrow \oint \mathbf{A} \cdot d\ell \equiv \Phi = n\Phi_0$$

trapped flux
↑
↑


$h/2e$
“superconducting”

flux quantum

$n = 0$ is trivial (no vortex!) and $|n| \geq 2$ is unstable, so restrict
 consideration to $n = (\pm)1$.

Since field extends over $\sim \lambda(T)$ into bulk metal,
 field at core ($\equiv H_0$) $\sim \Phi_0/\lambda^2(T)$.

Energetics of single vortex line (per unit length)



(1) “intrinsic” energies:

(a) field energy $\sim \frac{1}{2} \mu_0 H_0^2 \lambda^2 \sim \Phi_0^2 / (\mu_0 \lambda^2)$

(b) (minus) condensation energy $\sim -\frac{1}{2} \mu_0 H_c^2 \xi^2$

(c) flow energy: $v_s \sim \frac{\hbar}{2m} \frac{1}{r}$, for $r \leq \lambda(T)$, so

$$(c) \sim |\Psi_\infty|^2 \left(\frac{\hbar}{2m} \right)^2 \int_{\xi}^{\lambda} \frac{r dr}{r^2} \sim |\Psi_\infty|^2 \frac{\hbar^2}{(2m)^2} \ln(\lambda/\xi)$$

but $\lambda(T) = (\mu_0 |\Psi_\infty|^2 e^2 / m)^{-1/2}$, so

$$(c) \sim \left(\Phi_0^2 / \mu_0 \lambda^2(T) \right) \ln(\lambda/\xi)$$

dominant over
(a) and (b) for $\lambda(T) \gg \xi(T)$

Thus “intrinsic” energy per unit length of vortex lines for $\lambda \ll \xi$ is

$$E_0(T) \sim \left(\Phi_0^2 / (\mu_0 \lambda^2(T)) \cdot \ln(\lambda/\xi) \right).$$

- (2) On the other hand, the “extrinsic” energy saving due to admission of the external field is $\sim \mu_0 H_{ext} H_0 \sim H_{ext} (\Phi_0 / \lambda^2)$. Hence the condition for it to be energetically advantageous to admit a single vortex line is roughly

$$H_{ext} \sim (\Phi_0 / \lambda^2(T)) \cdot \ell n \kappa \quad (\kappa \equiv \lambda(T) / \xi(T) \neq f(T))$$

This defines the **lower critical field** H_{c1} .

What is the maximum field the superconductor can tolerate before switching to the N phase?

Roughly, defined by the point at which the vortex cores (area $\sim \xi^2(T)$) start to overlap. Since for near-complete penetration we must have $n\Phi_0 \sim H_{ext}$, this gives the condition

number of vortices/unit area

$$H_{ext} \sim (\Phi_0 / \xi^2(T))$$

This defines the **upper critical field** H_{c2}

Note that for $\lambda(T) \lesssim \xi(T)$, we have $H_{c1} \gtrsim H_{c2}$ and would expect type-I behavior.

Results of more quantitative treatment:

(a) condition for type-II behavior is $\kappa \equiv (\lambda(T)/\xi(T)) > 2^{-1/2}$

(b) in extreme type-II limit $\kappa \gg 1$,

$$H_{c1}(T) = (\Phi_0/4\pi\lambda^2(T))\ell n\kappa$$

(c) in same limit,

$$H_{c2}(T) = \Phi_0/2\pi\xi^2(T).$$

Note that quite generally we have up to logarithmic factors

$$H_{c1}(T) \cdot H_{c2}(T) \sim H_c^2(T)$$

└──────────┬ thermodynamic
critical field

hence if $H_c(T)$ held fixed, H_{c1} varies inversely to H_{c2}

Application: dirty superconductors

Alloying does not change $E_{cond}(T)$, hence $H_c^2(T)$, much. However, it drastically increases $\lambda(T)$ ($\xi(T)$ is also increased, but less). Hence κ is increased, and many elements which on type-I when pure became type-II when alloyed. As alloying increases, H_{c2} increases while H_{c1} decreases.

Summary of lecture 4

Ginzburg-Landau theory is special case of Landau-Lifshitz theory of 2nd order phase transitions: quantitatively valid only near T_c . Introduces order parameter (“macroscopic wave function”) $\Psi(\mathbf{r})$ which couples to vector potential $\mathbf{A}(\mathbf{r})$ with charge $2e$:

$$F\{\Psi(\mathbf{r})\} = \text{const.} - \alpha |\Psi(\mathbf{r})|^2 + \frac{1}{2} \beta |\Psi(\mathbf{r})|^4 + \\ \text{const.} \left| \left(\nabla - \frac{2ie}{\hbar} \mathbf{A}(\mathbf{r}) \right) \Psi(\mathbf{r}) \right|^2 + \frac{1}{2} \mu_0^{-1} (\nabla \times \mathbf{A}(\mathbf{r}))^2$$

2 characteristic lengths:

healing length $\xi(T)$

penetration depth $\lambda(T)$

both $\propto (T_c - T)^{-1/2}$ for $T \rightarrow T_c$.

Magnetic behavior depends on ratio $\kappa = \lambda(T)/\xi(T)|_{T \rightarrow T_c}$

For $\kappa > 2^{-1/2}$, (“type-I”) field completely expelled up to a thermodynamic critical field $H_c(T)$, at which point turns fully normal.

For $\kappa < 2^{-1/2}$, (“type-II”) field starts to penetrate at H_c in form of vortices, continues to do so up to H_{c2} where turns completely normal.