

SHANGHAI JIAO TONG UNIVERSITY
LECTURE 7
2017

Anthony J. Leggett

Department of Physics

University of Illinois at Urbana-Champaign, USA

and

Director, Center for Complex Physics

Shanghai Jiao Tong University



Recap: at $T = 0$ the structure of the MBWF is

$$\Psi = \prod_k \Phi_k \quad (\mathbf{k} \equiv (\mathbf{k} \uparrow, -\mathbf{k} \downarrow))$$

$$\Phi_k = u_k |00\rangle_k + v_k |11\rangle_k$$

and the specific values of u_k and v_k were found by minimizing $\langle \hat{H} - \mu \hat{N} \rangle$.

For $T \neq 0$ we expect intuitively that the description of the many-body system can still be factored into a product of descriptions of the occupation of the individual pair states $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$: technically

$$\hat{\rho} = \prod_k \hat{\rho}_k \quad \leftarrow \text{density matrix}$$

but now (a) all 4 occupation states will be realized with some probability

(b) quantities like Δ will be T -dependent

(c) at some $T_c \sim \Delta(T = 0)/k_B$ the collective bound state will cease to exist.



Recall: for given $\mathbf{k} \equiv (\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$ 4 occupational states

$$\begin{array}{cccc} |00\rangle, & |11\rangle, & |01\rangle, & |10\rangle \\ GP & EP & BP_1 & BP_2 \end{array}$$

and ground state has

$$\Delta \hat{x} \quad \psi_k = u_k |00\rangle + v_k |11\rangle \text{ corresponding to } \sigma_k \parallel \mathcal{H}_k = -\epsilon_k \hat{z} +$$

with an “energy” $-E_k \equiv |\mathcal{H}_k| \equiv (\epsilon_k^2 + |\Delta|^2)^{1/2}$. The limit $\Delta \rightarrow 0$ corresponds to the normal GS, and then $E_k \rightarrow |\epsilon_k|$. So the energy of the “ground pair” state relative to the normal ground state is

$$E_{GP} = |\epsilon_k| - E_k.$$

The EP (“excited pair”) state is formed by simply reversing the pseudospin k , so that

$$\psi_{k,EP} = v_k^* |00\rangle_k - u_k |11\rangle_k \quad (\text{orthogonal to } \psi_{k,GF})$$

This evidently costs an energy $2E_k$, so

$$E_{EP} = |\epsilon_k| + E_k$$

What about the BP (“broken pair”) states $BP_{1,2}$? These each correspond (relative to the N ground state) to kinetic energy (KE) $|\epsilon_k|$ and zero PE (no partner to scatter!), hence

$$E_{BP_{1,2}} = |\epsilon_k|$$

Thus the relative energies of the various states are

$$E_{BP} - E_{GP} = E_k, \quad E_{EP} - E_{GP} = 2E_k$$

State $BP_1(BP_2)$ has “Bogoliubov quasiparticle” in state $\mathbf{k} \uparrow (-\mathbf{k} \downarrow)$; state EP has quasiparticles in both $\mathbf{k} \uparrow$ and $\mathbf{k} \downarrow$ (hence $E_{EP} = 2E_{BP}$). (\Uparrow : but EP is really an “excitation of the condensate” whereas $BP_{1,2}$ are not).

Population of states: since all 4 states distinguishable, simple MB-Gibbs statistics applies, i.e. $P_n \propto \exp - \beta E_n$. Thus (taking E_{GP} as zero of E)

$$P_{GP} = Z^{-1}, P_{BP1} = P_{BP2} = Z^{-1} \exp - \beta E_k, P_{EP} = Z^{-1} \exp - 2\beta E_k$$

$$(E_k \equiv E_k(T))$$

$$Z = 1 + 2 \exp - \beta E_k + \exp - 2\beta E_k$$

A quantity of special interest is

$$F_k(T) \equiv \frac{1}{2} \langle \sigma_{xk} \rangle (T) = (\Delta(T)/2E_k)(P_{GP} - P_{EP})$$

$$= (\Delta(T)/2E_k(T)) \tanh \beta E_k(T)/2$$

Putting this into the equation

$$\Delta(T) = -V_0 \sum_k F_k(T)$$

we find

$$\Delta(T) = -V_0 \sum_k (\Delta(T)/2E_k(T) \tanh \beta E_k(T)/2)$$

or in the more general case ($V_0 \rightarrow V_{kk^1}$)

$$\Delta_k(T) = - \sum_{k'} V_{kk'} (\Delta_{k'}(T)/2E_{k'}(T)) \tanh \beta E_{k'}(T)/2$$



Finite-temperature BCS gap equation

As T increases from 0, $\Delta(T)$ decreases from $\Delta(0)$ to zero at a temperature T_c given by the linearized equation

$$\Delta_k(T_c) = - \sum_{k'} (V_{kk'} \Delta_{k'}(T_c) / 2 |E_{k'}|) \tanh \beta_c |E_{k'}| / 2 \quad (\beta_c \equiv 1/k_B T_c)$$

For the BCS contact potential ($V_{kk'} \rightarrow V_0$) this yields

$$[N(0)V_0]^{-1} = \int_0^{\epsilon_c} \frac{\tanh \beta \epsilon / 2}{\epsilon} d\epsilon = \ln(1.14 \beta_c \epsilon_c)$$

so comparing this with zero- T gap equation

$$[N(0)V_0]^{-1} = \ln(2\epsilon_c / \Delta(T = 0))$$

we have

$$\Delta(T = 0) = 1.76 k_B T_c$$

reasonably well satisfied for most “classical” superconductors

Examination of the gap equation at arbitrary $T < T_c$ shows that it is a function only of T/T_c

$$\Delta(T) = 1.76 k_B T_c f(T/T_c)$$

$$\text{with } f(z) \cong (1 - z^4)^{1/2}$$

(so for $T \rightarrow T_c$, $\Delta(T) \propto (1 - T/T_c)^{1/2}$)

Properties of a BCS superconductor at non zero T .

A. Condensate:

As we saw, the (F.T. of the) condensate wave function has the form at $T \neq 0$

$$F_k(T) = (\Delta(T)/2E_k(T)) \tanh \beta E_k(T)/2$$

so, in the wave function

$$F(\mathbf{r}) = \sum_k F_k \exp i\mathbf{k} \cdot \mathbf{r}$$

$$\equiv N(0) \int d\epsilon_k \frac{\sin kr}{kr} \frac{\Delta(T)}{(\epsilon_k^2 + \Delta^2(T))^{1/2}} \tanh \beta (\epsilon_k^2 + \Delta^2)^{1/2} / 2$$

the low energy cutoff (which determines the long distance behavior) gradually changes from $\sim \Delta(T = 0)$ to $\sim k_B T$. Since for $T \lesssim T_c$ these are of same order of magnitude, we have approximately

$$F(\mathbf{r}; T) \cong \Delta(T) \cdot N(0) \frac{\sin k_F r}{k_F r} \exp - r/\xi'(T)$$

where $\xi'(T) \sim \xi'(0)$. *i. e.*,

Cooper-pair radius is not sharply T -dependent
(in particular, does not diverge for $T \rightarrow T_c$ from below).

The number of Cooper pairs,

$$N_c(T) \int |F(\mathbf{r}; T)|^2 d\mathbf{r}$$

is proportional to $\Delta^2(T)$, hence for $T \rightarrow T_c$

$$N_c(T) \propto (1 - T/T_c)$$

Condensate is very “inert”, e.g. cannot be spin-polarized or (usually) flow in a way determined by walls. This applies both to GP and EP states (both have $S = 0$, COM momentum = 0). Hence such responses **determined entirely by BP states**. However, response is not simply proportional to the probability of occupation of BP states:

Ex: Pauli spin susceptibility

In field \mathcal{H} , $\Delta E = -\mu_B \mathcal{H} \sum_i S_i^z$. Hence, does not affect $|00\rangle$ or $|11\rangle$, but
↑
real spin not pseudospin!

shifts energies of BP states,

$$\Delta E_{BP_1} = -\mu_B \mathcal{H}, \quad \Delta E_{BP_2} = +\mu_B \mathcal{H}$$

Hence:

$$P_{BP_1} = \exp -\beta(E_k - \mu_B \mathcal{H}), \quad P_{BP_2} = \exp -\beta(E_k + \mu_B \mathcal{H})$$

and

$$\begin{aligned} \langle M_z \rangle &\equiv \mu_B \langle S_z \rangle \\ &= \mu_B^2 \sum_k (Z_k^{-1}) (\exp -\beta(E_k - \mu_B \mathcal{H}) - \exp -\beta(E_k + \mu_B \mathcal{H})). \end{aligned}$$

$$\text{with } Z_k(\mathcal{H}) = Z_k(0) + o(\mathcal{H}^2)$$



For $\mu_B \mathcal{H} \ll k_B T, \Delta(T)$ this gives

$$\langle M_z \rangle = \mu_B^2 \mathcal{H} \sum_k \frac{d}{dE_k} (\exp - \beta E_k) / Z_k = \mu_B^2 \mathcal{H} \beta \sum_k \text{sech}^2 \beta E_k / 2$$

and so


$$\chi \equiv \langle M_z \rangle / \mathcal{H} = \mu_B^2 \left(\frac{dn}{d\epsilon} \right) \beta \int_0^\infty \text{sech}^2(\beta E / 2) d\epsilon$$

In the normal state ($E \rightarrow \epsilon$) this correctly gives $\chi = \mu_B^2 dn/d\epsilon$, so

$$\chi(T) / \chi_n = \beta \int_0^\infty \text{sech}^2(\beta E(T) / 2) d\epsilon$$

↑
"Yosida function"

Note: Reason argument is relatively simple is that energy eigenstates ($\mathbf{k} \uparrow$) and ($-\mathbf{k} \downarrow$) carry a spin $+1/2$ ($-1/2$) respectively



The normal density

The “normal density” is defined as the **fraction of the electrons which can respond to a (transverse) static vector potential, in following sense:**

In presence of vector potential $\mathbf{A}(\mathbf{r})$

$$\mathbf{p} \rightarrow \mathbf{p} - e\mathbf{A}(\mathbf{r})$$

So KE becomes

$$\sum_i (\hat{p}_i - e\mathbf{A}(\mathbf{r}_i))^2 / 2m \equiv \sum_i \left(\frac{\hat{p}_i^2}{2m} - \frac{e}{m} \hat{p}_i \cdot \mathbf{A} + \frac{A^2(\mathbf{r}_i)}{m} \right)$$

↑
(ignore the order of operators)

and the current density $\mathbf{j}(\mathbf{r})$ is

$$\mathbf{j}(\mathbf{r}) = \frac{1}{2} \sum_i (\delta(\mathbf{r} - \mathbf{r}_i) (\hat{p}_i - e\mathbf{A}(\mathbf{r}_i)) / m + H.C.)$$

We already saw that the explicit term in $\mathbf{A}(\mathbf{r}_i)$ gives rise in the S phase, to the Meissner effect. But in the normal phase it is cancelled by the response of \hat{p}_i to the perturbation $\mathbf{p}_i \cdot \mathbf{A}(\mathbf{r}_i)$.

$$(\delta j / \delta A)_{pert} = + \frac{Ne^2}{m}$$

So: in S phase at $0 < T < T_c$ what is perturbative response of \mathbf{p} to \mathbf{A} ?

(almost) exact analogy to calculation of spin susceptibility:

$|00\rangle$ and $|11\rangle$ have total $\mathbf{P} = 0$, so cannot respond

$|10\rangle$ has momentum $\mathbf{p} = \hbar\mathbf{k}$, $|01\rangle$ has $\mathbf{p} = -\hbar\mathbf{k}$. Hence

$$\Delta E_{BP_1} = -e\hbar\mathbf{k} \cdot \mathbf{A}/m$$

$$\Delta E_{BP_2} = +e\hbar\mathbf{k} \cdot \mathbf{A}/m$$

Total induced momentum is

$$\mathbf{P} = \sum_{\mathbf{k}} \hbar\mathbf{k} (Z_{\mathbf{k}}^{-1}) \left(\exp - \beta \left(E_{\mathbf{k}} - \frac{e\hbar\mathbf{k} \cdot \mathbf{A}}{m} \right) - \exp - \beta \left(E_{\mathbf{k}} + \frac{e\hbar\mathbf{k} \cdot \mathbf{A}}{m} \right) \right)$$

and for $\hbar\mathbf{k} \cdot \mathbf{A} \ll k_B T, \Delta(T)$ this reduces to

$$J \equiv e \frac{\mathbf{P}}{m} \cong e^2 \hbar^2 \frac{k_F^2}{3m} \mathbf{A} \sum_{\mathbf{k}} (Z_{\mathbf{k}}^{-1}) \frac{d}{dE_{\mathbf{k}}} \exp - \beta E_{\mathbf{k}} \cong e^2 \frac{p_F^2}{3m} \beta \sum_{\mathbf{k}} \operatorname{sech}^2(\beta E_{\mathbf{k}}/2) \cdot \mathbf{A}$$

↑
directional averaging

In N state ($E \rightarrow \epsilon$) this correctly reduces to Ne^2/m , so ratio (“ ρ_n/ρ ”) of response in S state at temperature T to N –state value is

$$\rho_n/\rho = \beta \int_0^{\infty} (\operatorname{sech}^2 \beta E/2) d\epsilon$$

↑
Yosida function

↑ χ and ρ_n/ρ are untypically simple, because energy eigenstates are also eigenstates of σ and \mathbf{p} .



Summary of lecture 7

At $T \neq 0$ the BCS description is still a product over the different pair states $\mathbf{k} \equiv |\mathbf{k} \uparrow, -\mathbf{k} \downarrow\rangle$, but now all four states

$$|GP\rangle \equiv u_k |00\rangle + v_k |11\rangle$$

$$|BP1\rangle \equiv |10\rangle$$

$$|BP2\rangle \equiv |01\rangle$$

$$|EP\rangle \equiv v_k^* |00\rangle - u_k |11\rangle$$

are populated, and u_k and v_k are functions of T . The relative energies of the 4 states are

$$\left. \begin{aligned} E_{BP}(T) - E_{GP}(T) &= E_k(T) \\ E_{EP}(T) - E_{GP}(T) &= 2E_k(T) \end{aligned} \right\} E_k(T) \equiv (\epsilon_k^2 + |\Delta_k(T)|^2)^{1/2}$$

The self-consistent equation for the gap is

$$\Delta_k(T) = - \sum_{k'} V_{kk'} (\Delta_{k'}(T)/2E_{k'}(T)) \tanh(\beta E_{k'}(T)/2)$$

and has a nontrivial ($\Delta_k \neq 0$) solution only for $T < T_c$, where

$$k_B T_c = \Delta(T=0)/1.76$$

Condensate wave function $F(r; T)$ not strongly T -dependent:
no. of Cooper pairs $N_c(T) \sim \Delta^2(T)$, near $T_c \sim (1 - T/T_c)$
“Normal component” is essentially BP states: contributes to
“simple” quantities ($\chi, P_n \dots$) an amount $Y(T)$, e.g.

$$\chi(T)/\chi_n = Y(T) \equiv \beta \int_0^\beta \text{sech}^2(\beta E(T)/2) d\epsilon$$